arXiv:1304.6287v1 [gr-qc] 23 Apr 2013

Preprint typeset in JHEP style - HYPER VERSION

DCPT-13/15

Time dependent black holes and scalar hair

Sarah Chadburn[1]*, **Ruth Gregory**[1,2]†

[1] *Centre for Particle Theory, South Road, Durham, DH1 3LE, UK*

[2] *Perimeter Institute, 31 Caroline Street North, Waterloo, ON, N2L 2Y5, Canada*

ABSTRACT: We show how to correctly account for scalar accretion onto black holes in scalar field models of dark energy by a consistent expansion in terms of a slow roll parameter. We find an exact solution for the scalar field within our Hubble volume which is regular on both black hole and cosmological event horizons, and compute the back reaction of the scalar on the black hole, calculating the resulting expansion of the black hole. Our results are independent of the relative size of black hole and cosmological event horizons. We comment on the implications for more general black hole accretion, and the no hair theorems.

KEYWORDS: Black holes, scalar fields, accretion.

---

*Email: s.e.chadburn@durham.ac.uk

†Email: r.a.w.gregory@durham.ac.uk

Black holes are one of the most fascinating objects in classical general relativity. They represent the endpoint of gravitational collapse of a large mass object, independent of initial conditions. The prototypical black hole, the Schwarzschild solution, was first presented barely a year after Einstein put forward his new theory of gravity, yet it took half a century before relativists were confident of the interpretation of this solution, and began to construct a rigorous set of theorems describing the properties of black holes. One of the more poetically labelled set – the "no hair" theorems [1] – has been perhaps the most contentious. Originally demonstrated for static vacuum black holes, [2, 3], the no hair theorems have been refined and extended to a wide range of interacting particle and gravitational theories, [4, 5, 6]. Nonetheless, there are many known "violations" of the no hair theorems, from the dressing of black holes by scalar or gauge condensates [7], to literal hair, in the form of topological defects which extend out from the black hole to infinity, [8].

The issue of the most basic hair however, scalar hair, is thought to be understood, yet is perhaps the most perplexing. For a wide range of potentials, it is easy to show that a static black hole cannot have a nontrivial scalar field on its horizon (unless the spacetime is asymptotically anti-de Sitter, [9], or the scalar potential has been specifically tailored, [10]). However, within cosmology, scalar fields are widely used, not only for inflation, but as an expedient model of quintessence, [11], or late time acceleration (see [12] for a review of dark energy models). Indeed, the standard exponential scalar potential - widely used to give power law acceleration, was shown to be in conflict with spherically symmetric asymptotically flat or de Sitter static black hole spacetimes, [13]. How then can these cosmological scalars interact with a black hole horizon? Generally, the scalar field is varying on a very slow time scale relative to the light crossing time of the black hole, thus we might expect that it has little effect on the horizon itself, yet, like the inexorable flow of time, the scalar must evolve cosmologically, and it seems contradictory that it is pinned to a single value for all time at the black hole.

In fact, the resolution of the conflict is quite straightforward: The no hair theorems were explored in the context of static black holes, and quite clearly a rolling cosmological scalar can never be static. Correspondingly, the presence of a black hole in a cosmology implies that the cosmology cannot be spatially homogeneous, and therefore does not have the canonical FRW form. While we can use cosmological perturbation theory to describe the far field effect of the black hole, we cannot describe the full spatial geometry including the event horizon within the context of perturbation theory – and it is the effect of the event horizon which is critical in determining the behaviour of the scalar. We therefore need a time-dependent black hole solution.

Exact black hole solutions are surprisingly thin on the ground, given how ubiquitous black hole are in nature, and time dependent ones even more so. The first dynamical black hole 'solution' was postulated by McVittie, [15], who modified the

Schwarzschild solution to include an FRW-like cosmology, with the black hole horizon at fixed comoving radius and a fluid energy density dependent only on time. However, as pointed out in [16], one expects the black hole to break the spatial homogeneity of the cosmology, and thus (except in the special case of a de Sitter universe) this unrealistic set-up leads to a singularity on the would be horizon. The Vaidya solution, [14], also time dependent, is somewhat special, as the mass of the black hole now becomes dependent on a null coordinate, requiring a null matter source. In cosmology however, we expect a black hole to accrete general stress-energy, and as already noted, solutions in the literature with nontrivial scalar fields have very specialized potentials, with fixed mass black holes. Thus a direct approach to finding an exact solution seems to have been unsuccessful in that solutions do not account for an evolution of the black hole horizon area due to scalar accretion.

An alternate approach in the literature is to find a probe solution for the scalar field, i.e. one in which the scalar field evolves on a Schwarzschild background, [17, 18]. Here, near the horizon, the scalar field must be a function of the advanced null coordinate, $v \sim e^{t+r^*}$, which parametrises the black hole future event horizon. One can then estimate the back-reaction on the event horizon using the energy momentum of this approximate solution. A problem with this approach however, is that the coordinates being used are local static Schwarzschild coordinates, which do not correspond to the cosmological flow of time at large distances and therefore give no guarantee that any 'solution' will be well behaved at cosmological horizons. One can also worry that estimates using local intuitive notions of energy, rather than an analysis of the Einstein equations with a scalar source, might be misleading.

In order to be confident that probe calculations give a good estimate of scalar accretion onto black holes, we require a time-dependent scalar field with a time dependent black hole. Here, we present a resolution of this problem, by expanding the equations of motion for the geometry and the scalar field order by order in a "slow roll" parameter. We show how to put a time dependent scalar on the black hole, and how this extends to the cosmological solution at large distances. Our solution is fully nonperturbative in the radial direction, thus is valid independent of the relative sizes of the black hole and cosmological horizons. We also present the back reaction of the scalar field on the black hole, and calculate the expansion rate of the black hole due to scalar accretion. An advantage of our method is that we can identify the event horizons accurately as null surfaces, therefore the usual ambiguity of apparent vs. event horizons does not occur.

To motivate our approach, consider a universe in which acceleration is driven by a scalar field with an exponential potential $W(\phi) = M^4 e^{-\beta\phi}$. Writing the solution in conformal time gives

$$ds^2 = \left[\frac{\eta_0}{\eta}\right]^{2+\frac{2\varepsilon}{1-\varepsilon}} \left[(d\eta - d\rho)(d\eta + d\rho) - \rho^2 d\Omega_{\rm II}^2\right] \tag{1}$$

$$\phi = \phi_0 - \frac{2\varepsilon}{\beta(1-\varepsilon)} \ln \eta \tag{2}$$

where $\varepsilon = \beta^2/2\kappa = \frac{M_p^2 W'(\phi)^2}{2W(\phi)^2}$ is none other than the slow roll parameter for the potential W, and represents the power of the acceleration. Clearly, if we assume that $\varepsilon \ll 1$, $[\eta_0/\eta]^{\frac{2\varepsilon}{1-\varepsilon}} \sim 1 - 2\varepsilon \ln |\eta/\eta_0|$, and hence this metric is perturbatively close to the de Sitter (dS) metric, over a Hubble time interval. From (2), $\phi = \phi_0 + \mathcal{O}\left(\varepsilon/\kappa\right)^{1/2}$, and thus our cosmology is roughly dS, with a small correction of order $\mathcal{O}(\varepsilon^{1/2})$ for the scalar, and $\mathcal{O}(\varepsilon)$ for the geometry.

Now consider adding a black hole to this cosmological rolling scalar solution. The geometry will not now have constant curvature spatial slices, but we do expect an SO(3) symmetry, corresponding to the spherical symmetry of the black hole. We also expect both time and radial dependence in the metric. The general metric may be written in the form [19, 20]

$$ds^2 = 4e^{2\nu} B^{-1/2} dU dV - B d\Omega_{\rm II}^2. \tag{3}$$

This form of the metric elucidates both the symmetry of the spacetime (SO(3)), as well as the remaining gauge freedom (the conformal group in the U, V directions). By rewriting the coordinates in lightcone form, it will be clear how to deal with the horizons present in the anticipated solution (the cosmological and black hole event horizons), as well as how to change gauge to analytically extend across these horizons. In particular, the null coordinates allow us to identify the actual event horizons of the solution, as opposed to apparent horizons, as an horizon is of course always a null surface, defined by U or $V =$ constant.

The coupled black hole scalar equations we have to solve are

$$B_{,UV} = 2\left(\kappa W(\phi) B^{1/2} - B^{-1/2}\right) e^{2\nu} \tag{4}$$

$$\nu_{,UV} = \frac{1}{2}\left(\kappa W(\phi) B^{-1/2} + B^{-3/2}\right) e^{2\nu} - \frac{\kappa}{2}\phi_{,U}\phi_{,V} \tag{5}$$

$$\phi_{,UV} = -W_{,\phi}(\phi) B^{-1/2} e^{2\nu} - \frac{1}{2B}\left(B_{,U}\phi_{,V} + B_{,V}\phi_{,U}\right) \tag{6}$$

$$B_{,VV} = 2\nu_{,V} B_{,V} - \kappa B \phi_{,V}^2 \tag{7}$$

$$B_{,UU} = 2\nu_{,U} B_{,U} - \kappa B \phi_{,U}^2 \tag{8}$$

For a constant scalar field (as one might naively suppose from the no hair theorems) the Einstein equations have Schwarzschild de Sitter (SdS) as a general solution, [19]. To briefly review, if ϕ is constant, (7) and (8) can be integrated directly to show that B must be a function of $[F(V) + G(U)]$, where F and G are arbitrary integration functions, and $e^{2\nu} = F'(V)G'(U)B'(F+G)$. Equation (4) then gives

$$B' = \frac{4}{3}\kappa W_0 B^{3/2} - 4B^{1/2} + 8GM = -4\sqrt{B} N(\sqrt{B}) \tag{9}$$

where $8GM$ is an integration constant (suggestively labeled!), and we see that the function N defined above is none other than the SdS potential, with $\kappa W_0/3 = H^2$, the Hubble parameter. One can then change coordinates to $r = B^{1/2}$, $t = 2(G(U) - F(V))$, to obtain

$$ds^2 = \frac{F'G'B'}{B^{1/2}} 4dUdV - Bd\Omega^2_{\rm II} = N(r)dt^2 - \frac{dr^2}{N(r)} - r^2 d\Omega^2_{\rm II} \tag{10}$$

i.e. the Schwarzschild de Sitter metric in static coordinates.

Although the static coordinates are useful to identify horizons in the SdS metric, they are distinct from the "cosmological" coordinates, in which the metric would at large r asymptote a standard cosmological de Sitter spacetime; they also have the disadvantage that they are singular at both the black hole, and cosmological, event horizons. Writing r_i ($r_N < 0 \leq r_h < r_c$) as the roots of $N(r)$, and $R_i = r_i/(1 - 3H^2 r_i^2) = 1/N'(r_i)$, we have a locally regular null coordinate system at each horizon given by

$$\begin{cases} V = R_c \exp\left[-2F/R_c\right] \ , \ \ U = R_c \exp\left[2G/R_c\right] & \text{CEH} \\ v = R_h \exp\left[-2F/R_h\right] \ , \ \ u = -R_h \exp\left[-2G/R_h\right] & \text{BH} \end{cases} \tag{11}$$

the latter being the standard Kruskal coordinates for the SdS black hole, and the former a set of Kruskals for the cosmological event horizon, provided we identify $F = -(t + r^*)/4$, and $G = (t - r^*)/4$, with r^* being the usual tortoise coordinate.

Now suppose that ϕ is not constant, but slowly rolling, then we can solve our spacetime and scalar field order by order in the slow roll parameter. The solution will be perturbative in ε, but, crucially, *nonperturbative* in the radial coordinate across our Hubble volume. Writing $\phi = \phi_0 + 2\varepsilon\phi_1/\beta$ in (6) gives

$$\phi_{1,UV} = 3H^2 B^{-1/2} B'F'G' - \frac{B'}{2B}\left(G'\phi_{1,V} + F'\phi_{1,U}\right) \tag{12}$$

It is reasonably clear that the form

$$\phi_1 = 2\lambda(G(U) - F(V)) + \varphi(F + G) \tag{13}$$

will satisfy (12) provided $[B\varphi']' = 3H^2 B^{1/2} B'$, hence

$$\varphi = -\sum_i \left[H^2 r_i + \frac{C}{r_i^2}\right] R_i \ln|r - r_i| + \frac{C}{2GM}\ln r \tag{14}$$

with C an integration constant.

For a nonsingular solution, the ϕ field must be regular in a locally regular coordinate system at both the black hole (r_h), and cosmological (r_c), future event horizons, thus we get two constraints on λ and C, solved by

$$C = -H^2 \frac{r_h^2 r_c^2 (r_h + r_c)}{r_h^2 + r_c^2} \quad , \quad \lambda = \frac{(r_c - r_h)}{r_h^2 + r_c^2} \tag{15}$$

Pulling this together we can write the ϕ field in the static SdS coordinates:

$$\begin{aligned}\phi =& \phi_0 + \frac{2\varepsilon}{\beta}\lambda\Big[t - R_c \ln|r-r_c| + R_h \ln(r-r_h) \\ & - \frac{R_N}{\lambda}\left(H^2 r_N + \frac{C}{r_N^2}\right)\ln(r+r_c+r_h) - \frac{r_h r_c \ln r}{(r_c-r_h)}\Big]\end{aligned} \tag{16}$$

which is manifestly nonsingular at the horizons, and plotted in figure 1.



Figure 1: Contours of constant ϕ (thick blue) within our Hubble volume. For comparison, contours of constant t and r are shown in grey, and a constant *cosmological* time contour $(U+V)$ shown in red. The horizons are labelled, as are the singularity and future infinity.

It is useful to note the asymptotic forms of the scalar field. Near the cosmological event horizon

$$\phi \sim \phi_0 + \frac{2\varepsilon}{\beta} 2\lambda R_c \ln(U+V) \tag{17}$$

which gives the correct cosmological rolling scalar solution to order $\mathcal{O}(\varepsilon)$ as $r_h \to 0$, (cf (2)). On the other hand, near the event horizon,

$$\phi \sim \phi_0 + \frac{2\varepsilon}{\beta} 2\lambda R_h \ln v \tag{18}$$

in agreement with the linearized solutions of [17, 18]. Clearly, ϕ is regular from the black hole event horizon out to the cosmological event horizon, with regular derivatives, and thus its energy momentum is finite in this region. We can therefore compute the backreaction on the geometry to get a consistent solution to order $\mathcal{O}(\varepsilon)$.

Writing $B = r^2(1+\varepsilon\delta_1)$ and $\nu = \nu_{SdS}+\varepsilon\delta_2$, the procedure follows the integrability argument for the SdS solution: First integrate (7) and (8) to obtain two expressions for δ_2

$$2\delta_2 = \delta_1 + \frac{rV\delta_{1,V}}{2R_cN} + g(U) + I_V = \delta_1 - \frac{rU\delta_{1,U}}{2R_cN} + f(V) + I_U \tag{19}$$

where

$$\begin{aligned} I_V &= 2\int \frac{B\phi_{1,V}^2}{B_{,V}}dV = \int \frac{rdr}{N^2}(H^2r + \frac{C}{r^2} - \lambda)^2 \ , \\ I_U &= 2\int \frac{B\phi_{1,U}^2}{B_{,U}}dU = \int \frac{rdr}{N^2}(H^2r + \frac{C}{r^2} + \lambda)^2 \ . \end{aligned} \tag{20}$$

Next, consistency of (19) determines the general form of δ_1:

$$\frac{-r\delta_1}{R_cN} = \ln(UV)(I_V - I_U) - \int \frac{2f}{V}dV + \int \frac{2g}{U}dU + h(r) \tag{21}$$

with (4) giving a second order ODE for $h(r)$, solved by

$$h(r) = \int \frac{dr}{rN^2R_c}\left[h_1 - \int [12H^2r^2\varphi + (rN)'\left(I_U + I_V\right)]\right] \tag{22}$$

Regularity at both horizons then uniquely determines the various integration constants (or functions) and hence $\delta B, \delta\nu$.

The general expressions for δB and $\delta\nu$ are rather cumbersome, but by focussing on the event horizons, it is straightforward to obtain results of most physical interest. For the cosmological event horizon, the $\{U,V\}$ coordinate system is appropriate, with the CEH being at $V = 0$, and parametrized by U. Since $r =$ const. along the horizons, (8) gives

$$B = r_c^2\left(1 - 8\varepsilon\lambda^2R_c^2\ln|U|\right) \simeq r_c^2|U|^{-8\varepsilon\lambda^2R_c^2} \tag{23}$$

Setting $r_h \to 0$ gives $4\lambda^2R_c^2 = 1$, and hence $B \propto U^{-2\varepsilon}$, in complete agreement with the cosmological horizon area of the pure rolling scalar solution, (1).

Of more interest however is the accretion of scalar field onto the black hole. Here, using the Kruskal $\{u, v\}$ system and (7), we get

$$B = r_h^2\left(1 + 8\varepsilon\lambda^2R_h^2\ln v\right) \simeq r_h^2 v^{8\varepsilon\lambda^2R_h^2} \tag{24}$$

We can compare this with an order of magnitude estimate based on naive physical notions of mass and energy flow, [17, 18]. The flow of energy into the black hole should be governed by the difference of the energy momentum tensor from being null, $T_0^0 - T_r^r$, which is of order $\dot{\phi}^2 \sim W'^2/H^2 \sim \varepsilon H^2/\kappa$. Integrating this over the horizon gives $\kappa\delta\dot{M} \sim r_h^2\varepsilon H^2$, or using the relation between horizon radius and mass: $\delta A \sim r_h^3\varepsilon H^2\delta t$. Of course, on the horizon, t is not a good coordinate, and we should instead use advanced time: $t \to 2r_h\ln v$, which gives $\delta A \sim A\varepsilon(\delta\ln v)r_h^2/r_c^2$, in agreement with (24).

For an astrophysical black hole, this accretion rate is glacially slow, and far outweighed by the local environment, in which the accretion disc far outweighs local interstellar matter, let alone this cosmologically coasting scalar. However, the fact that naive local estimates of the backreaction of accretion of scalar matter in this set-up are fully backed up by this radially nonperturbative analytic calculation means we should have confidence in these physically motivated techniques.

The other feature we should emphasise about our results is precisely their non-perturbative nature in the radial coordinate. While the expansion is perturbative in time, our solution for the scalar, and hence the geometry, is exact across the space-time between the black hole and cosmological event horizons. Thus, we can apply these results to black holes whose event horizon constitutes a significant fraction of the Hubble volume. These black holes accrete at a similar rate to the increase in area of the cosmological horizon.

We should point out that while we have a time dependent scalar field in a time dependent cosmological black hole background, this should not be viewed as a violation of the no hair theorems. The solution corresponds to a rolling cosmological scalar field in which there is a black hole, and the rolling of the scalar adjusts to its presence. The scalar does roll in the vicinity of the black hole, although it lags behind the cosmological evolution, in the sense that the constant ϕ contours lie in front of the constant η contours in figure 1. However, the black hole does not have any scalar charge – there is no 1-parameter solution for ϕ in the black hole background. The solution (13) is not the most general solution, there will be wave like fluctuations around this background, but we have not been able to find a family of solutions with spacelike dependence, which would be a signature of a scalar charge on the black hole. Thus, our results should be viewed as a way of reconciling the "no hair" intuition with more general time dependent situations.

Finally, to our knowledge, this is the first analytic spacetime presented with a time dependent accreting black hole that results from consideration of a physically realistic matter system (rather than reverse engineering from a given metric). The results apply to a general potential, and only require that the scalar field is slowly rolling. As such, they represent a testing ground for investigation of black hole phenomena in the time dependent regime.

Acknowledgments

We would like to thank Anne Davis for helpful discussions. RG is supported in part by STFC (Consolidated Grant ST/J000426/1), in part by the Wolfson Foundation and Royal Society, and in part by Perimeter Institute for Theoretical Physics. SC is supported by an EPSRC studentship. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Economic Development and Innovation.

References

[1] R. Ruffini and J. A. Wheeler, Physics Today **24**, 30 (1971).

[2] B. Carter, Phys. Rev. Lett. **26**, 331 (1971).

[3] J. D. Bekenstein, Phys. Rev. D **5**, 1239 (1972).

[4] S. L. Adler and R. B. Pearson, Phys. Rev. D **18**, 2798 (1978).

[5] J. D. Bekenstein, Phys. Rev. D **51**, 6608 (1995).

[6] A. E. Mayo and J. D. Bekenstein, Phys. Rev. D **54**, 5059 (1996) [gr-qc/9602057].

[7] R. Bartnik and J. Mckinnon, Phys. Rev. Lett. **61**, 141 (1988).

[8] A. Achucarro, R. Gregory and K. Kuijken, Phys. Rev. D **52**, 5729 (1995) [gr-qc/9505039].

[9] T. Torii, K. Maeda and M. Narita, Phys. Rev. D **64**, 044007 (2001).

[10] K. G. Zloshchastiev, Phys. Rev. Lett. **94**, 121101 (2005) [hep-th/0408163].

[11] R. R. Caldwell, R. Dave and P. J. Steinhardt, Phys. Rev. Lett. **80**, 1582 (1998) [astro-ph/9708069].

[12] E. J. Copeland, M. Sami and S. Tsujikawa, Int. J. Mod. Phys. D **15**, 1753 (2006) [hep-th/0603057].

[13] S. J. Poletti and D. L. Wiltshire, Phys. Rev. D **50**, 7260 (1994) [Erratum-ibid. D **52**, 3753 (1995)] [gr-qc/9407021].

[14] P. Vaidya, Proc. Indian Acad. Sci. A **33**, 264 (1951).

[15] G. C. McVittie, Mon. Not. Roy. Astron. Soc. **93**, 325 (1933).

[16] N. Kaloper, M. Kleban and D. Martin, Phys. Rev. D **81**, 104044 (2010) [arXiv:1003.4777 [hep-th]].

[17] T. Jacobson, Phys. Rev. Lett. **83**, 2699 (1999) [astro-ph/9905303].

[18] A. V. Frolov and L. Kofman, JCAP **0305**, 009 (2003) [hep-th/0212327].

[19] P. Bowcock, C. Charmousis and R. Gregory, Class. Quant. Grav. **17**, 4745 (2000) [hep-th/0007177].

[20] C. Charmousis and R. Gregory, Class. Quant. Grav. **21**, 527 (2004) [gr-qc/0306069].